

Qorshe Hawleedka Xasaasiyad Ba'an "Anaphylaxis"

Dadka u baahan daawaynta xaalada degdeg ah ee EPINEPHRINE

"Anaphylaxis waxaa weeye waxyeelo xasaasiyad leh taas oo si degdeg ah u bilaabanta lagana yaabo inay dhimasho sababto."

(Maxadka Xasaasiyada Qaranka iyo Cudurada Neefsashada ku faafa, 2010)

Sawir

Magac: _____ Taariikhda Dhalashada: _____

WUXUU XASAASIYAD KU LEEYAHAY: _____

Macluumaad ku saabsan Xiiqda: Haa (Xasaasiga daran ee halista badan ku sugan) Maya

Ma qaadan karaa dawada: Haa Maya Ma maamuli karaa daawada: Haa Maya

Qiyaasta Daawada

Qiyaasta EPINEPHRINE:

Ilaa 55 Rodol. (25 kiloo) **Ka badab 55 Rodol.** (25 kiloo)

EpiPen Jr. (0.15 miligaraam) EpiPen (0.3 miligaraam)

Adrenaclick (0.15 miligaraam) Adrenaclick (0.3 miligaraam)

*Nooca Antihistamine + Qiyaasta daawada:

Benadryl (sidoo kale loo yaqaano Diphenhydramine)

12.5 miligaraam (1 mulqaacada shaaha lagu qaado ama 1 la calalin karo)

25 miligaraam (2 mulqaacadood oo shaaha lagu qaado ama 2 la calalin karo)

50 miligaraam (4 mulqaacadood oo shaaha lagu qaado ama 4 la calalin karo)

Antihistamine Kale: _____

Waxyeelo aada u leh cunnooyinka soo socda: _____

SIDAAS DARAADDEED:

Haddii baadhitaan lagu sameeyey, si degdeg ah u sii EPINEPHRINE haddiba ay jiraan ASTAAMO XANUUN OO KASTA haddii cunnada xasaasiyada keenaysa ay suurtoagal tahay in la cuno.

Haddii baadhitaan lagu sameeyey, si degdeg ah u sii EPINEPHRINE haddii cunnada xasaasiyada keenaysa aanu shaki ku jirin in la cunay, xaataa haddii aan la oggaan astaamo xanuun.

ASTAMIHII DARAN EE XANUUN kasta kadib marka laga shakiyo ama loo yaqaano dheefshiidis:

Mid ama inkabandan oo kamida kuwa soo socda:

Sanbab: Neefsi kooban, hinraagid, qufac soo noqnoqda

Wadne: Midab khafiifa, niyad jab, miyir doorsoon, garaaca wadnaha oo tamardaran, dawakhaad, asqoobid

Cunaha: Tigtigan, xabeeb, neefsiga oo ku dhiba/liqida oo ku dhibta

Afka: Barar is xidha (carabka iyo/ama dabnaha)

Dubka: Jidhka gebi ahaantii oo casuus iyo cuncun yeesha

Ama **isku jir** astaamaha xanuun oo ka yimaada qaybo kaladuwan oo jidhka ah:

Dubka: Jidh casuus iyo cuncun yeesha, finan cuncun leh, barar (indhaha, dabnaha)

Midhicirka: Matag, xanuun casiraad wata

1. ISLA MARKIIBA KU MUD EPINEPHRINE

2. Wac telefoon lambarka 911

3. Bilow inaad la socoto (sida lagu faahfaahiyey xagga hoose)

4. Sii daawayn dheeraada: *

- Antihistamine
- Inhaler (bronchodilator) haddii uu xiiq leeyahay

*Antihistamines iyo inhalers/ bronchodilators looma isticmaalo in lagu daaweeyo waxyeelada daran (anaphylaxis). ISTICMAAL EPINEPHRINE.

ASTAAMO XANUUNKA CAADIGA AH ee kaliya:

Afka: Af cuncun leh

Dubka: Waxoogaa Jidh casuus iyo cuncun leh oo afka/wajiga ku gadaaman, cuncun caadi ah

Midhicirka: Yalaalugo Caadi ah/Xanuun

1. SII ANTIHISTAMINE

2. La joogista ardayga; heegan gali khabirada daryeelka caafimaadku iyo waalidka/masuulka

3. Haddii astaamaha xanuunku ay isa soo taraan (eeg xagga sare) ISTICMAAL EPINEPHRINE

4. Bilow inaad la socoto (sida lagu faahfaahiyey xagga hoose)

Xaaladaha gaarka ah: _____

Dabagal joogtaynta

QIYAASTA LABAAD EE DAAWO QAADASHADA ee EPINEPHRINE waxaa la siin karaa 5 daqiiqadood ama wax kabadan kadib haddii astaamaha xanuunku ay sii socdaan ama ay dib u soo ifbaxaan.

La joogista qofka; heegan galinta khabirada daryeelka caafimaadka iyo waalidka/masuulka. U sheeg in EPINEPHRINE la siiyey. Waxaad oggaataa waqtigii EPINEPHRINE la siiyey. Waxyeelada daran, waxaad ka fikirtaa inaad qofka dhabarka u jiiifso lugahana aad kor u taagto. Daawee qofka xaataa haddii aan waalidka la heli karin. Waxaad eegtaa dariiqada dhabarka danbe/ee ku lifaaqan irbada iskeed isku wada.

Saxiixa Biyaha: _____

Telefoon

Taariikh

Magaca Daabacan: _____

Saxiixa Waalid/Masuul: _____

Telefoon


Taariikh

**EPIPEN® Auto-Injector iyo
EPIPEN Jr® Dariiqooyinka Auto-Injector**

- Marka hore, kasaar EPIPEN Auto-Injector bacda ay ku jirto



- Soo jiid daboolka leh midabka cirkoo kale ee laga furo
- Si adag u xaji qaybta sare ee u dhaw (had iyo jeer ku qabo bawdyaha)



- Horay iyo diba u dhufo oo si adag ugu cadaadi caarada sare ee ku beegan bawdada oogadeeda. Ku xaji bawdada qiyaas ahaan 10 ilbidhiqsi. Ka saar EPIPEN Auto-Injector oo ku duug jiida ilaa 10 daqiiqadood wax kabadan



DEY® iyo DEY logo, EpiPen®, EpiPen 2-Pak®, iyo EpiPen Jr 2-Pak waa astaan ganacsi oo u sajjilan Dey Pharma, L.P.

**Adrenallick™ 0.3 miligaraam iyo
Adrenallick™ 0.15 miligaraam Dariiqooyinka**



Ka saar furarka MIDABKA CAWLAN ee ay ku calaamadaysan "1" iyo "2".

Dhig caarada sare wareegsan ee Midabka CASAANKA AH meel ku beegan bawdada oogadeeda sare, si adag hoos ugu cadaadi ilaa iyo inta ay irbadu galayso. Ku xaji ilaa 10 ilbidhiqsi, kadibna ka saar.



Xidhmada cunnooyinka xasaasiyada lihi waxaa haboon inay ku jiraan ugu yaraan laba qiyaas daawo qaadashada epinephrine, daawooyinka kale sida uu u sheegay dhaqtarka ardaygu, iyo nuqulkan shaqo hawleedka Cunnada Xasaasiyada.

Xidhmada qalabka waa in la raaciyaa ardayga haddii uu/ay ka maqan yahay/ka maqan tahay garoonada dugsiga (tusaale ahaan, socdaalada shaqada).

La xidhiidhitaanka

Wac telefoon lambarka 911 (Rescue squad: _____) Dhakhtar: _____ Telefoon: _____

Waalid /Masuul: _____ Telefoon: _____

Xidhiidhitaanada Kale ee Xaaladaha Degdega ah

Magac/Xidhiidh: _____ Telefoon: _____

Magac/Xidhiidh: _____ Telefoon: _____